

A characterization of a class of maximum nonlinear functions

Doreen Hertel and Alexander Pott
 Institute for Algebra and Geometry
 Otto-von-Guericke-University Magdeburg
 D-39016 Magdeburg

February 2, 2008

Abstract

Maximum nonlinear functions $F : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_{2^m}$ are widely used in cryptography because the coordinate functions $F_\beta(x) := \text{tr}(\beta F(x))$, $\beta \in \mathbb{F}_{2^m}^*$, have large distance to linear functions. More precisely, the Hamming distance to the characteristic functions of hyperplanes is large. One class of maximum nonlinear functions are the Gold power functions x^{2^k+1} , $\gcd(k, m) = 1$. We characterize these functions in terms of the distance of their coordinate functions to characteristic functions of subspaces of codimension 2 in $\mathbb{F}_{2^m}$.

Keywords. m -sequence, maximum nonlinear function, Gold power function, Walsh transform

1 Introduction

The finite field with 2^m elements is denoted by $\mathbb{F}_{2^m}$. The multiplicative group of the field is denoted by $\mathbb{F}_{2^m}^*$. We may also view $\mathbb{F}_{2^m}$ as an m -dimensional vector space over $\mathbb{F}_2$. The trace function is the linear mapping $\text{tr} : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$ defined by $\text{tr}(x) = \sum_{i=0}^{m-1} x^{2^i}$. It is well known that the mappings tr_β defined by $\text{tr}_\beta(x) = \text{tr}(\beta x)$ are linear, again, and all 2^m linear mappings $\mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$ can be represented like this. The reader is referred to [16] for background from the theory of finite fields.

The **Hamming distance** between two **boolean** functions $f, g : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$ is the number of elements $x \in \mathbb{F}_{2^m}$ such that $f(x) \neq g(x)$. The distance d between f

and the linear function tr_γ is $2^{m-1} - \frac{t_\gamma}{2}$, where

$$(1) \quad t_\gamma = \sum_{x \in \mathbb{F}_{2^m}} (-1)^{tr(\gamma x + f(x))}.$$

This is easily seen since $t_\gamma = 2^m - 2d$. The distance d to the affine function $x \mapsto tr_\gamma(x) + 1$ is $2^{m-1} + \frac{t_\gamma}{2}$ (now $-t_\gamma = 2^m - 2d$). We say that f is highly nonlinear if the smallest distance to all (affine) linear functions is very high. In other words, the maximum value of t_γ (where $\gamma \in \mathbb{F}_{2^m}$) is small. We denote the maximum value for $|t_\gamma|$ the **linearity** of f :

$$(2) \quad \mathcal{L}(f) := \max_{\gamma \in \mathbb{F}_{2^m}} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{tr(\gamma x + f(x))} \right|.$$

It is well known that

$$(3) \quad \mathcal{L}(f) \geq 2^{\frac{m}{2}}$$

where equality may occur (of course, only for m even), see [5], for instance. A function f satisfying $\mathcal{L}(f) = 2^{\frac{m}{2}}$ is called a **bent function**. They exist for every even m , see [5]. Bent functions are basically the same objects as certain difference sets, see [14] and [1]. Note that $\mathcal{L}(f) = 2^m$ if f is linear or affine.

Many authors formulate this problem in a slightly different (though equivalent) way: They define the **nonlinearity** of f to be

$$(4) \quad \mathcal{N}(f) := 2^{m-1} - \frac{1}{2} \mathcal{L}(f).$$

This number is the smallest Hamming distance between f and the set of all (affine) linear functions $tr(\gamma x)$ and $tr(\gamma x) + 1$ (which is called the 1st order Reed-Muller code). The number $\mathcal{N}(f)$ is the so called covering radius of this code.

The goal is to maximize $\mathcal{N}(f)$. This maximum is known precisely only in the case m even, hence the covering radius problem is solved if m is even. If m is odd, the problem seems to be much harder.

The numbers t_γ are sometimes called the **Walsh coefficients** $\mathcal{W}_f(\gamma)$. If $f(x) = x^d$, we simply write $\mathcal{W}_d(\gamma)$. In this paper, we do not need the full power of the Walsh transformation, therefore we just refer to the literature for more information about this important concept, see [9], for instance. However, we use the notation $\mathcal{W}_d(\gamma)$ to denote (1) for the power mapping x^d . Moreover we mention that the function f is uniquely determined by its Walsh spectrum

$$\{\mathcal{W}_f(\gamma) : \gamma \in \mathbb{F}_{2^m}\}.$$

This also holds if $f : \mathbb{F}_{2^m} \rightarrow \mathbb{C}$ is a complex-valued function, and we define

$$\mathcal{W}_f(\gamma) := \sum_{x \in \mathbb{F}_{2^m}} f(x) (-1)^{tr(\gamma x)}.$$

We may interpret the integers $\mathcal{W}_d(\gamma)$ also in terms of the intersection between certain sets. First of all note that any function $f : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$ defines a subset D_f of $\mathbb{F}_{2^m}$:

$$D_f := \{x \in \mathbb{F}_{2^m} : f(x) = 1\}.$$

Vice versa, every subset gives rise to a mapping $\mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$.

Specifically, we define for d with $\gcd(d, 2^m - 1) = 1$:

$$\begin{aligned} D_d &:= \{x \in \mathbb{F}_{2^m} : \text{tr}(x^d) = 1\} \\ H^0(\alpha) &= \{x \in \mathbb{F}_{2^m} : \text{tr}(\alpha x) = 0\} \\ H^1(\alpha) &= \{x \in \mathbb{F}_{2^m} : \text{tr}(\alpha x) = 1\}. \end{aligned}$$

If $\alpha \neq 0$, the sets $H^0(\alpha)$ and $H^1(\alpha)$ are subspaces of codimension 1 in $\mathbb{F}_{2^m}$ (hyperplanes), i.e. they have size 2^{m-1} . Therefore, we obtain for $\alpha \neq 0$

$$\begin{aligned} \mathcal{W}_d(\alpha) &= 2^m - 2(|D_d \cap H^0(\alpha)| + |(\mathbb{F}_{2^m} \setminus D_d) \cap (\mathbb{F}_{2^m} \setminus H^0(\alpha))|) \\ (5) \quad &= 2^m - 4|D_d \cap H^0(\alpha)| \end{aligned}$$

and

$$(6) \quad -\mathcal{W}_d(\alpha) = 2^m - 4|D_d \cap H^1(\alpha)|.$$

Since $|D_d| = 2^{m-1}$ we have $\mathcal{W}_d(0) = 0$.

Now we turn our attention to **vectorial** functions $F : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_{2^m}$.

Let $F : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_{2^m}$ be arbitrary. We consider the **coordinate functions** $F_\beta(x) := \text{tr}(\beta \cdot F(x))$ from $\mathbb{F}_{2^m}$ to $\mathbb{F}_2$. The smallest nonlinearity of all nonzero coordinate functions of F is called, similar to the boolean case, the **nonlinearity** of F :

$$\mathcal{N}(F) = \min_{\beta \in \mathbb{F}_{2^m}^*} \mathcal{N}(F_\beta).$$

Similarly, the linearity is

$$\begin{aligned} \mathcal{L}(F) &= \max_{\beta \in \mathbb{F}_{2^m}^*} \mathcal{L}(F_\beta) \\ &= \max_{\beta, \gamma \in \mathbb{F}_{2^m}, \beta \neq 0} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\gamma \cdot x + \beta \cdot F(x))} \right| \end{aligned}$$

and the connection between theses two numbers is, like in (4),

$$(7) \quad \mathcal{N}(F) = 2^{m-1} - \frac{1}{2}\mathcal{L}(F).$$

Similar to the case of boolean functions f , we have a (rather easy to prove) lower bound

$$\mathcal{L}(F) \geq 2^{\frac{m+1}{2}}$$

where equality may occur if m is odd, see again [5], for instance. Functions with $\mathcal{L}(F) = 2^{\frac{m+1}{2}}$ are called **maximum nonlinear** or **almost bent**. In this case, we know the linearities of all the coordinate functions:

$$(8) \quad \sum_{x \in \mathbb{F}_{2^m}} |(-1)^{\text{tr}(\gamma x + \beta F(x))}| \in \{0, \pm 2^{\frac{m+1}{2}}\},$$

see [5], again. Also the multiplicities are known. The following table shows how often the sum in (8) (for a fixed $\beta \neq 0$) takes the three different values:

$$(9) \quad \begin{array}{cc} \text{value in (8)} & \text{multiplicity} \\ \hline 0 & 2^{m-1} \\ 2^{\frac{m+1}{2}} & 2^{m-2} \pm 2^{\frac{m-3}{2}} \\ -2^{\frac{m+1}{2}} & 2^{m-2} \mp 2^{\frac{m-3}{2}}. \end{array}$$

More precisely, if $\text{tr}(\beta F(0)) = 0$ then $2^{\frac{m+1}{2}}$ occurs $2^{m-2} + 2^{\frac{m-3}{2}}$ times, otherwise it occurs $2^{m-2} - 2^{\frac{m-3}{2}}$ times.

Let k be an integer with $\gcd(k, m) = 1$. Then the mappings $\mathbb{F}_{2^m} \rightarrow \mathbb{F}_{2^m}$ defined by

$$x^{2^k+1} \quad \text{or} \quad x^{2^{2k}-2^k+1}$$

are maximum nonlinear. The first examples are called the **Gold** power mappings, see [8], the second class of mappings are the **Kasami** power mappings, see [9].

There are two more classes of maximum nonlinear functions known (**Welch** and **Niho** case), and they are also power mappings x^d . They have been proved to be maximum nonlinear only recently, see [3] and [11].

If $\gcd(k, m) \neq 1$ or if m is even, the linearities of the power mappings x^{2^k+1} and $x^{2^{2k}-2^k+1}$ are also known, provided that $\gcd(d, 2^m - 1) = 1$, see [9].

It is interesting to know that, up to now, all maximum nonlinear mappings can be constructed from these four classes. As usual, functions that can be constructed from each other using some specified procedure are called equivalent. In the case of functions, there are different ways to define *equivalence*. A nice way to unify these is contained in [4]. We refer the reader to [2] where it is shown that the equivalence described in [4] is indeed more general than the *classical* affine equivalence. When we say that all known maximum nonlinear functions can be constructed from each other, we do not mean that they are all *affine equivalent*, but that they can be constructed from each other according to Proposition 3 in [4], see [2].

In this paper we consider maximum nonlinear power functions x^d . Perhaps, this class contains more mappings than just described. In view of the connection between the Walsh coefficients of $\text{tr}(\beta x^d)$ and the intersection between D_d and

$H^i(\alpha)$ in (5) and (6), we obtain the following intersection numbers between $H^0(\alpha)$, $\alpha \in \mathbb{F}_{2^m}^*$, and D_d :

$$(10) \quad \begin{array}{cc} |D_d \cap H^0(\alpha)| & \text{multiplicity} \\ \hline 2^{m-2} & 2^{m-1} \\ 2^{m-2} - 2^{\frac{m-3}{2}} & 2^{m-2} + 2^{\frac{m-3}{2}} \\ 2^{m-2} + 2^{\frac{m-3}{2}} & 2^{m-2} - 2^{\frac{m-3}{2}}. \end{array}$$

In this paper, we consider the intersection between D_d and subspaces of codimension 2. We can characterize the Gold power mappings in terms of these intersection sizes.

If F is a maximum nonlinear power mapping x^d , then one can show that x^d has to be a permutation, i.e. $\gcd(d, 2^m - 1) = 1$. It seems that this argument did not yet appear in the literature. It is actually due to Dobbertin, and it will appear in [15]. The proof uses the fact that any maximum nonlinear mapping is an almost perfect nonlinear function. We emphasize that not all maximum nonlinear functions are bijective: Just by adding a suitable linear function αx one gets nonbijective functions.

In the case of power mappings with $\gcd(d, 2^m - 1) = 1$, the computation of the (non)linearity simplifies. In (11), put $\beta = \eta^d$ which is possible since $\gcd(d, 2^m - 1) = 1$. In (12), we replace γ by $\alpha\eta$. Finally in (13) we note that ηx runs through $\mathbb{F}_{2^m}$ if x does:

$$(11) \quad \mathcal{L}(x^d) = \max_{\gamma, \beta \in \mathbb{F}_{2^m}, \beta \neq 0} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\gamma \cdot x + \beta \cdot x^d)} \right|$$

$$(12) \quad = \max_{\gamma, \eta \in \mathbb{F}_{2^m}, \eta \neq 0} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\gamma \cdot x + (\eta \cdot x)^d)} \right|$$

$$(13) \quad = \max_{\alpha, \eta \in \mathbb{F}_{2^m}, \eta \neq 0} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\alpha\eta \cdot x + (\eta \cdot x)^d)} \right|$$

$$(14) \quad = \max_{\alpha \in \mathbb{F}_{2^m}} \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\alpha x + x^d)} \right|.$$

Note that

$$(14) \quad \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\alpha x + x^d)} = -1 + \sum_{x \in \mathbb{F}_{2^m}^*} (-1)^{\text{tr}(\alpha x + x^d)}.$$

This observation implies some connections between the linearity of power mappings and the crosscorrelation between m -sequences and their decimations, as we will describe next.

Binary Sequences $a = (a_i)_{i \geq 0}$ ($a_i \in \{0, 1\}$) are called **periodic** with period n if $a_i = a_{i+n}$ for all i . The **autocorrelation** of a binary sequence a with period n is defined by

$$c_t(a) := \sum_{i=0}^{n-1} (-1)^{a_i + a_{i+t}}.$$

The integer t is called a **phase shift** of the sequence a . Since the sequence is n -periodic, we may compute the indices modulo n .

A sequence with n odd and $c_t(a) = -1$ for all $1 \leq t \leq n-1$ is called **perfect**, see [12] for more background on perfect sequences.

Let ζ be a primitive element of $\mathbb{F}_{2^m}$. The sequences $a = (a_i)$ with $a_i = \text{tr}(\zeta^i)$ are called **m -sequences**. They have period $2^m - 1$ and they are perfect. Other classes of perfect sequences are known. We refer the reader to the chapter on difference sets in [1] since perfect sequences correspond to a certain class of cyclic difference sets, see also [12].

Similarly to the autocorrelation, we define the **crosscorrelation** between two binary sequences a and b of period n by

$$c_t(a, b) := \sum_{i=0}^{n-1} (-1)^{a_i + b_{i+t}}.$$

Finally, we define the **d -decimation** $a^{[d]}$ of an n -periodic sequence $a = (a_i)$ by $a_i^{[d]} := a_{id}$. Note that $a^{[d]}$ is an m -sequence corresponding to the primitive element ζ^d if a is the m -sequence defined by $a_i := \text{tr}(\zeta^i)$ and $\gcd(d, 2^m - 1) = 1$.

If ζ is a primitive element of $\mathbb{F}_{2^m}$, we may reformulate the righthand side of (14):

$$\sum_{x \in \mathbb{F}_{2^m}^*} (-1)^{\text{tr}(\alpha x + x^d)} = \sum_{i=0}^{2^m-2} (-1)^{\text{tr}(\alpha \zeta^i + \zeta^{id})} = \sum_{i=0}^{2^m-2} (-1)^{\text{tr}(\zeta^{i+t} + \zeta^{id})}$$

(define t by $\alpha = \zeta^t$). This shows that the linearity of the power mapping x^d is the same as -1 plus the maximum crosscorrelation value between an m -sequence and its d -decimation.

2 Main Theorem

Theorem 1 *Let m be odd and let x^d be a maximum nonlinear power function on $\mathbb{F}_{2^m}$. Let*

$$H^{i,j}(\alpha, \beta) := \{x : \text{tr}(\alpha x) = i, \text{tr}(\beta x) = j\}.$$

Then $d = 2^k + 1$ for some integer k with $\gcd(k, m) = 1$ (i.e. d is a Gold exponent) if and only if

$$(15) \quad |H^{i,j}(\alpha, \beta) \cap D_d| \in \{2^{m-3}, 2^{m-3} \pm 2^{\frac{m-3}{2}}\}$$

for all $\alpha, \beta \in \mathbb{F}_{2^m}^*$, $\alpha \neq \beta$, and $i, j \in \mathbb{F}_2$.

The sets $H^{i,j}(\alpha, \beta)$, $\alpha \neq \beta$, $\alpha, \beta \in \mathbb{F}_{2^m}^*$, are precisely the subspaces of dimension $m - 2$ in $\mathbb{F}_{2^m}$. The set D_d has some interesting properties. It is the set of 2^{m-1} points in the m -dimensional vector space $\mathbb{F}_{2^m}$ over $\mathbb{F}_2$. If d is a Gold exponent, this set is a non-degenerate quadric, see [9], for instance. If m is odd, there is up to equivalence only one non-degenerate quadric in $\mathbb{F}_{2^m}$, and the intersection between this quadric and subspaces of codimension 2 must be the three values described in (15). This is well known to geometers, see [10], for instance. It follows from [7] that the only quadrics corresponding to the coordinate functions of maximum nonlinear power mappings are nondegenerate: In [7], the intersection sizes between quadrics Q and hyperplanes are determined whenever $|Q| = 2^{m-1}$. This applies to the situation of maximum nonlinear power mappings x^d and m odd, since in this case $|D_d| = 2^{m-1}$ (because x^d is bijective). It turns out that the intersection sizes in 10 occur only in the nondegenerate case.

It is natural to ask whether there are values d such that D_d is not a nondegenerate quadric but has the same intersection sizes with hyperplanes. These objects are called by geometers **quasi-quadrics**. Many examples of quasi-quadrics are known, see [6], for instance. Note that all maximum nonlinear power mappings yield quasi-quadrics. Our research was motivated by the question whether the quasi-quadrics constructed from maximum nonlinear functions may also behave like quadrics if the intersection sizes with subspaces of codimension 2 are considered. The answer, given by Theorem 1, is no.

An interesting corollary is the following:

Corollary 2 *The only maximum nonlinear power mappings x^d on $\mathbb{F}_{2^m}$ such that D_d is a quadric are the Gold power mappings.*

Before we are going to prove our Theorem, let us mention the following Proposition which may be of interest in its own:

Proposition 3 *Let x^d be a maximum nonlinear power mapping on $\mathbb{F}_{2^m}$ with $\gcd(d, m) = 1$. Then*

$$(16) \quad |H^{i,j}(\alpha, \beta) \cap D_d| \in \{2^{m-3} + h \cdot 2^{\frac{m-5}{2}} : -3 \leq h \leq 3\},$$

where $\alpha, \beta \in \mathbb{F}_{2^m}^*$, $\alpha \neq \beta$.

Proof. We define

$$S^{i,j}(\alpha, \beta) = |H^{i,j}(\alpha, \beta) \cap D_d|$$

and

$$S^i(\alpha) = |H^i(\alpha) \cap D_d|.$$

Assume $\alpha \neq \beta$, $\alpha, \beta \in \mathbb{F}_{2^m}^*$. We obtain

$$\begin{aligned} |\mathcal{W}_d(\alpha + \beta)| &= \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\alpha x + \beta x + x^d)} \right| \\ &= \left| \sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(\alpha x + i + \beta x + j + x^d)} \right| \\ &= 2^m - 2(3 \cdot 2^{m-1} - 2S^i(\alpha) - 2S^j(\beta) - 2 \cdot 2^{m-2} + 4S^{i,j}(\alpha, \beta)) \\ &= -2^m + 4S^i(\alpha) + 4S^j(\beta) - 8S^{i,j}(\alpha, \beta). \end{aligned}$$

Because of (5) and (6), we have

$$|\mathcal{W}_d(\alpha + \beta)| = 2^m \pm \mathcal{W}_d(\alpha) \pm \mathcal{W}_d(\beta) - 8S^{i,j}(\alpha, \beta),$$

hence

$$(17) \quad S^{i,j}(\alpha, \beta) = 2^{m-3} \pm \frac{1}{8}(\pm \mathcal{W}_d(\alpha + \beta) \pm \mathcal{W}_d(\alpha) \pm \mathcal{W}_d(\beta)).$$

This shows that there are only the seven possible values for $S^{i,j}(\alpha, \beta)$ stated in the Proposition. $\square$

The proof of Theorem 1 reduces to the proof of an interesting property of the trace function. This Theorem has been independently obtained by Ph. Langevin and P. Véron [15]. The proof given in their paper is different from ours. The Langevin-Véron proof is shorter and more elegant, though less elementary.

Theorem 4 is not true in the case m even. For instance, if we take $m = 8$ and $d = 51$, then $\text{tr}(x^d + (x + 1)^d + 1) = 0$ for all $x \in \mathbb{F}_{2^8}$. This example can be extracted from the proof, since line (8) of the following algorithm does not produce the desired element w if m is even. Note that $\gcd(51, 2^8 - 1) \neq 1$. If we restrict ourselves to the case $\gcd(d, 2^m - 1) = 1$, Theorem 4 remains true also if m is even, see Lemma 2 in [13].

Theorem 4 *Let m be odd and $d \in \{2, \dots, 2^m - 2\}$ odd. We have*

$$\text{tr}(x^d + (x + 1)^d + 1) = 0$$

for all $x \in \mathbb{F}_{2^m}$, if and only if $d = 2^k + 1$ for one $k \in \mathbb{N}$.

We postpone the proof of Theorem 4 to the next Section. We are now going to show that it is sufficient to prove Theorem 4 in order to check Theorem 1.

Let x^d be a maximum nonlinear power function on $\mathbb{F}_{2^m}$, hence the Walsh spectrum $\{\mathcal{W}_d(\alpha) | \alpha \in \mathbb{F}_{2^m}\}$ contains only the three value $\pm 2^{\frac{m+1}{2}}$ and 0. We define the function $b : \mathbb{F}_{2^m} \rightarrow \mathbb{F}_2$ by

$$b(\alpha) = \begin{cases} 1 & \text{if } \mathcal{W}_d(\alpha) \neq 0 \\ 0 & \text{otherwise.} \end{cases}$$

If all or precisely one of the values $\mathcal{W}_d(\alpha)$, $\mathcal{W}_d(\alpha)$ and $\mathcal{W}_d(\alpha)$ in equation (17) are $\neq 0$, it is impossible that $S^{i,j}(\alpha, \beta) \in \{2^{m-3}, 2^{m-3} \pm 2^{\frac{m-3}{2}}\}$. Therefore, $b(\alpha) + b(\beta) = b(\alpha + \beta)$, hence b is linear and therefore

$$b(x) = \text{tr}(\gamma x) \quad (= \text{tr}_\gamma(x))$$

for some $\gamma \in \mathbb{F}_{2^m}^*$. If we think of $\text{tr}(x)$ as an element in $\mathbb{C}$, we obtain

$$\begin{aligned} \mathcal{W}(\text{tr}_\gamma)(\omega) &= \sum_{x \in \mathbb{F}_{2^m}} \text{tr}(\gamma x) \cdot (-1)^{\text{tr}(\omega x)} \\ &= \sum_{x \in \mathbb{F}_{2^m}, \text{tr}(\gamma(x))=1} (-1)^{\text{tr}(\omega x)} \\ (18) \quad &= \begin{cases} -2^{m-1} & \text{if } \omega = \gamma \\ 2^{m-1} & \text{if } \omega = 0 \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

On the other hand, the function b satisfies

$$b(x) = \frac{1}{2^{m+1}} [\mathcal{W}_d(x)]^2.$$

We compute the Walsh transform again:

$$\begin{aligned} \mathcal{W}(b)(\omega) &= \sum_{x \in \mathbb{F}_{2^m}} \frac{1}{2^{m+1}} (\mathcal{W}_d(x))^2 (-1)^{\text{tr}(\omega x)} \\ &= \frac{1}{2^{m+1}} \sum_{x, y, z \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(x(y+z+\omega) + (y^d+z^d))} \\ &= \frac{1}{2^{m+1}} \sum_{y, z \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(y^d+z^d)} \underbrace{\sum_{x \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(x(y+z+\omega))}}_{= \begin{cases} 2^m & \text{if } z = \omega + y \\ 0 & \text{otherwise} \end{cases}} \\ &= \frac{1}{2} \sum_{y \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(y^d + (y+\omega)^d)}. \end{aligned}$$

We compare this with (18) and obtain

$$(19) \quad \sum_{y \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(y^d + (y+\omega)^d)} = \begin{cases} -2^m & \text{if } \omega = \gamma \\ 2^m & \text{if } \omega = 0 \\ 0 & \text{otherwise.} \end{cases}$$

The case $\omega = \gamma$ implies

$$(20) \quad \text{tr}(y^d + (y + \gamma)^d) = 1 \quad \text{for all } y \in \mathbb{F}_{2^m}.$$

We can show that necessarily $\gamma = 1$:

$$\begin{aligned} \text{tr}((y + \gamma)^d) &\stackrel{(20)}{=} \text{tr}(y^d) + 1 \\ &= \text{tr}(y^{2^l d}) + 1 \\ &\stackrel{(20)}{=} \text{tr}((y^{2^l} + \gamma)^d) \\ &= \text{tr}((y + \gamma^{2^{m-l}})^d) \end{aligned}$$

for all $l = 0, \dots, m-1$ and $y \in \mathbb{F}_{2^m}$. Suppose, that $\gamma \neq \gamma^{2^k}$ for some $k \in \mathbb{N}$, then $\text{tr}(y^d + (y + \gamma^{2^k})^d) = \text{tr}(y^d + (y + \gamma)^d) = 1$, thus $\sum_{y \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(y^d + (y + \gamma^{2^k})^d)} = \sum_{y \in \mathbb{F}_{2^m}} (-1)^{\text{tr}(y^d + (y + \gamma)^d)} = -2^m$. This is a contradiction to the uniqueness of γ . Thus we have $\gamma^{2^l} = \gamma$ for all $l = 0, \dots, m-1$, and therefore $\gamma = 1$.

Since m is odd we have $\text{tr}(1) = 1$. Therefore

$$(21) \quad \text{tr}(y^d + (y + 1)^d + 1) = 0$$

for all $y \in \mathbb{F}_{2^m}$. This shows that it is enough to prove Theorem 4.

3 Proof of Theorem 4

If d satisfies (21), then each $d' \in D$ with $D := \{2^i d \bmod (2^m - 1) : i = 0, \dots, m-1\}$ also satisfies (21). We choose the smallest odd d' in D , and from now on, we denote this element by d .

Let $w(a)$ be the binary weight of a . If $a = \sum_{i=0}^n z_i 2^i$ is the binary representation of a , we denote the vector $(z_n, \dots, z_0)$ by $\bar{a}$. We have $\sum_{i=0}^n z_i = w(a)$. In the following proof, all integers a that occur are $\leq 2^m - 1$, i.e. $\bar{a}$ is a vector of length at most m . By adding 0's, if necessary, we assume that $\bar{a}$ is always a vector of length m . Let $v = (v_{m-1}, \dots, v_0)$ be a vector of length m . We denote by $v^{(t)}$ the cyclic shift of the vector v about t positions to the left, i.e. $v^{(t)} = (v_{m-t-1}, \dots, v_0, v_{m-1}, \dots, v_{m-t})$.

Let $d' = 2^i d \bmod (2^m - 1)$, then $\bar{d}' = \bar{d}^{(t)}$, in particular $w(d') = w(d)$.

We define the polynomial p by

$$p(x) := x^d + (x + 1)^d + 1$$

and q by

$$q(x) := \text{tr}(p(x)) = \sum_{i=0}^{m-1} (p(x))^{2^i}.$$

We have $q(0) = 0$, therefore we have to show that

$$(22) \quad q(\alpha) = 0 \quad \text{for all } \alpha \in \mathbb{F}_{2^m}^*.$$

Let $T = \{t_1, \dots, t_n\}$ denote the set of exponents which occur in p . We define the set

$$T(t) = \{0 \leq s \leq 2^m - 2 : \bar{s}^{(i)} = \bar{t}, i = 0, \dots, m-1\}.$$

We obtain

$$q(x) = \sum_{t \in T} \sum_{s \in T(t)} x^s.$$

In order to prove (22), we have to show that every exponent occurs an even number of times in $q(x)$.

Let $\bar{d} = (z_{m-1}, \dots, z_0)$ be the binary vector corresponding to d . Since d is odd, we have $w(d) \neq 1$. If $d = 2^k + 1$ is a Gold exponent, then $w(d) = 2$ and $q(x)$ satisfies (22) (note that $p(x) = x^{2^k} + x$ in this case). Hence we may assume $w(d) \geq 3$.

If $w(d) = 3$, then $d = 2^k + 2^l + 1$ and $k > l > 0$. For the polynomials p and q we obtain

$$\begin{aligned} p(x) &= x^{2^k+2^l} + x^{2^k+1} + x^{2^l+1} + x^{2^k} + x^{2^l} + x \\ q(x) &= \sum_{i=0}^{m-1} \left((x^{2^k+2^l})^{2^i} + (x^{2^k+1})^{2^i} + (x^{2^l+1})^{2^i} + x^{2^i} \right). \end{aligned}$$

In $p(x)$, the exponents of binary weight 1 (resp. 2) occur three times, therefore we have an odd number of exponents of weight 1 (resp. 2) in $q(x)$, and therefore $q(x)$ cannot satisfy (22). This argument can be generalized: If $z = w(d)$ then there are precisely $\binom{z}{i}$ exponents t in $p(x)$ with $w(t) = i$, $1 \leq t \leq d-1$. Note that x^d and 1 do not occur in $p(x)$. If z is not a power of 2, at least one of these binomial coefficients is odd. Therefore, we only have to consider the case $z = 2^n$, $n > 1$.

Let v be a binary vector of length m . A **subvector** $w = (w_{m-1}, \dots, w_0)$ of v is a binary vector $w \neq 0, v$ of length m such that $v_i = 0$ implies $w_i = 0$. The set of all subvectors of $\bar{d}$ is the set of the binary vectors of the exponents that occur in $p(x)$. In order to show that (22) holds, we have to prove that the cardinality of the multiset

$$S(s) := \{ \bar{s}^{(i)} : \bar{s}^{(i)} \text{ subvector of } \bar{d}, 0 \leq i \leq m-1 \}$$

is even for all $s \in T$. Note that it is possible that $\bar{s}^{(i)} = \bar{s}^{(j)}$ for $i \neq j$.

We define a **gap** to be a substring v of the form $0\dots 0$. The number s of 0's in this substring is called the length of the gap, similarly for **runs** which are substrings of the form $1\dots 1$. More precisely: There is an i such that $z_i = z_{i+1} = \dots = z_{i+s-1} = 0$, where the indices are computed modulo m , i.e. we view $\bar{d}$ as a “cyclic” vector. If $v = (v_i v_{i+1} \dots v_j)$ is a substring, we say that the indices $i, \dots, j$ are contained in v .

By the following algorithm we construct a subvector w of $\bar{d}$ such that $|S(w)|$ is odd. Therefore, (22) is not satisfied.

Algorithm

Input: binary vector $\bar{d} = (z_{m-1}, \dots, z_0)$ of weight $z = 2^n, n \in \mathbb{N}, m$ odd

Output: subvector w of $\bar{d}$ such that $|S(w)|$ is odd

- (1) $l :=$ maximum length of a run in $\bar{d}$;
 $s :=$ multiplicity of a run of length l in $\bar{d}$;
 $v :=$ run of length l ;
 $s_{old} := m + 1; x_{old} := 0$;
- (2) while (w is not defined) do
- (3) $y := (y_{m-1}, \dots, y_0)$ with

$$y_i = \begin{cases} 1 & \text{if } i \text{ is contained in a substring } v \text{ and } z_i \text{ is } 1 \\ 0 & \text{otherwise.} \end{cases}$$
- (4) if $z \neq l \cdot s$ then $w := y$; end if;
- (5) if $z = l \cdot s$ then
 $x :=$ minimum length of a gap between two substrings v in y ;
 $L :=$ gap of length x ;
if $s = 1$ then
- (6) if $s_{old} = m + 1$ then $w := \bar{d} - (0\dots 010)$; end if;
- (7) if $s_{old} \neq m + 1$ then $w := (0\dots 0v_{old}L_{old}v_{old})$; end if;
- end if;
- (8) if $s = 2$ then $w := (0\dots 0vLv)$; end if;
- (9) if $s > 2$ then
 $s_{old} := s; l_{old} := l; x_{old} := x; L_{old} := L; v_{old} := v$;
let v denote a substring of type $(v_{old}Lv_{old}\dots Lv_{old})$ in $\bar{d}$
with maximum number l of 1's;
 $v := (v_{old}Lv_{old}\dots Lv_{old})$ with v_{old} occurs l/l_{old} -times;
 $s :=$ multiplicity of v in $\bar{d}$;
end if;
- end if;
- end while;

The algorithm terminates if $z \neq l \cdot s$ or $s \leq 2$. Note, if the case $z \neq l \cdot s$ does not

occur then such an s exists because $0 < s < s_{old}$ in each step of the algorithm.

Line (4): If $z \neq l \cdot s$, i.e. $y \neq \bar{d}$ and $w = y$ is a subvector of $\bar{d}$. We have $|S(w)| = 1$, because none of the cyclic shifts $w^{(t)} \neq w$ is a subvector of $\bar{d}$. Suppose the vector $w^{(t)}$ with $w^{(t)} \neq w$ is a subvector of $\bar{d}$. Note, that w and $w^{(t)}$ have the same number of 1. If $w^{(t)} \neq w$, then there exists a 1 in $\bar{d}$ and this 1 is in $w^{(t)}$ and not in w . Because $w^{(t)}$ is a cyclic shift of w , this 1 is in a string v , therefore this 1 is in w . This is a contradiction to the definition of w .

Line (5): If $z = l \cdot s$, then $l = 2^{l'}$ and $s = 2^{s'}$. We call the gaps L_j , $j = 1, \dots, s$ between the runs v . Now we know, that $\bar{d}$ has the form

$$\bar{d} = (L_s v L_{s-1} v \dots L_2 v L_1 v).$$

If $s > 1$, the number of gaps is even. Since m is odd, the number of gaps with odd length and the number of gaps with even length is odd. Therefore the maximum and minimum gap have different length. Note by the choice of $d \in D$ to be odd it follows that L_s is one of the maximum gaps and has length $> x$, the minimum length of a gap.

Line (6): If $z = l \cdot s$ with $s = 1$ and $s_{old} = m + 1$ then $l \geq 4$ and $\bar{d} = (0 \dots 01 \dots 1)$. For $w = \bar{d} - (0 \dots 010)$ we have $|S(w)| = 1$.

Line (7): If $z = l \cdot s$ with $s = 1$ and $s_{old} \neq m + 1$, then $s_{old} \geq 4$. The vector $\bar{d}$ has the form

$$\bar{d} = (L_s v) = (L_{s_{old}} v_{old} L v_{old} \dots L v_{old} L v_{old}),$$

where L is the gap of length x_{old} . We obtain $|S(w)| = s_{old} - 1$ is odd.

Line (8): If $s = 2$, then is $\bar{d} = (L_2 v L_1 v)$. The gap L_2 is longer than the gap L_1 . It is easy to see that $|S(w)| = 1$.

Line (9): The new initialisation for the next while loop. □

We illustrate the algorithm with an example. Here we have $m = 23$ and $d = 1 + 2^2 + 2^4 + 2^7 + 2^9 + 2^{11} + 2^{15} + 2^{17}$.

Input: $\bar{d} = (00000101000101010010101)$

(1) $z := 8; l := 1; s := 8; v := 1; s_{old} := 24; x_{old} := 0;$

(3) $y := \bar{d};$

(5) $x := 1; L := 0;$

(9) $s_{old} := 8; l_{old} := 1; x_{old} := 0; L_{old} := 0; v_{old} := 1;$

$$y = (00000\underline{101000101010010101});$$

$l := 3; v := 10101; s := 2;$

(3) $y := (00000000000v00v);$

(4) $w := y;$

Output: $w := (00000000000101010010101).$

References

- [1] T. BETH, D. JUNGnickel, AND H. LENZ, *Design Theory*, Cambridge University Press, Cambridge, 2 ed., 1999.
- [2] L. BUDAGHYAN, C. CARLET, AND A. POTT, *New constructions of almost bent and almost perfect nonlinear polynomials*, in Abstract Book of the International Workshop on Coding and Cryptography, Bergen (Norway), P. Charpin and Ø. Ytrehus, eds., 2005, pp. 306–315.
- [3] A. CANTEAUT, P. CHARPIN, AND H. DOBBERTIN, *Binary m -sequences with three-valued crosscorrelation: A proof of Welch's conjecture*, IEEE Trans. Inf. Theory, 46 (2000), pp. 4–8.
- [4] C. CARLET, P. CHARPIN, AND V. ZINOVIEV, *Codes, bent functions and permutations suitable for DES-like cryptosystems*, Des. Codes Cryptogr., 15 (1998), pp. 125–156.
- [5] F. CHABAUD AND S. VAUDENAY, *Links between differential and linear cryptanalysis*, in Advances in Cryptology – EUROCRYPT 94, A. D. Santis, ed., vol. 950 of Lecture Notes in Computer Science, New York, 1995, Springer-Verlag, pp. 356–365.
- [6] F. DE CLERCK, N. HAMILTON, C. M. O'KEEFE, AND T. PENTTILÄ, *Quasi-quadrics and related structures.*, Australas. J. Comb., 22 (2000), pp. 151–166.
- [7] R. A. GAMES, *The geometry of quadrics and correlations of sequences*, IEEE Trans. Inf. Th., 32 (1986), pp. 423–426.
- [8] R. GOLD, *Maximal recursive sequences with 3-valued recursive cross-correlation function*, IEEE Trans. Inf. Th., 14 (1968), pp. 154–156.
- [9] T. HELLESETH AND P. V. KUMAR, *Sequences with low correlation*, in Handbook of coding theory, Vol. I, II, North-Holland, Amsterdam, 1998, pp. 1065–1138.
- [10] J. HIRSCHFELD, *Projective geometries over finite fields. 2nd ed.*, Oxford Mathematical Monographs. Oxford: Clarendon Press., 1998.
- [11] H. H. HOLLMANN AND Q. XIANG, *A proof of the Welch and Niho conjectures on crosscorrelations of binary m -sequences*, Finite Fields Appl., 7 (2001), pp. 253–286.
- [12] D. JUNGnickel AND A. POTT, *Perfect and almost perfect sequences*, Discrete Applied Mathematics, 95 (1999), pp. 331–359.

- [13] G. M. KYUREGHYAN, *Differentially affine maps*, in Abstract Book of the International Workshop on Coding and Cryptography, Bergen (Norway), P. Charpin and Ø. Ytrehus, eds., 2005, pp. 296–305.
- [14] E. S. LANDER, *Symmetric designs: an algebraic approach*, vol. 74 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 1983.
- [15] P. LANGEVIN AND P. VÉRON, *On the nonlinearity of power functions*. to appear in Designs, Codes and Cryptography, 2005.
- [16] R. LIDL AND H. NIEDERREITER, *Finite Fields*, vol. 20 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2nd ed., 1997.